


Gynaecomastia



Correction of Enlarged Male Breasts

Plastic Surgery @ it's Best

درماپلاست
dermaplast

مستشفى الدكتور طارق
DR. TARIQ HOSPITAL

www.dermaplast.com.bh

Welcome to dermaplast

The single most important factor in the success of aesthetic (cosmetic) surgery is the surgeon you select and his team.

Dr. Tariq Saeed is the renowned Bahraini, Plastic, Aesthetic and Laser surgeon who has been in plastic surgery practice since 1983 and trained in Ireland, UK and the USA. Dr. Tariq holds fellowships in surgery from both the Royal College of Surgeons in Ireland and Edinburgh. He holds higher Fellowships in Plastic Surgery from Ireland and the University of Miami in the USA. Dr. Tariq is an assistant professor of Surgery at the Arabian Gulf University in Bahrain and is an external examiner for the Royal College of Surgeons in Ireland. Dr. Tariq is a founding member and past president of the GCC Association of Plastic Surgeons and also a founding member and the Past President of the Pan Arab Association of Burns & Plastic Surgery along with his membership of international medical societies including international membership of the American Society of Plastic Surgeons, American Society for Aesthetic Plastic Surgery, and International Society of Aesthetic Surgery.

Dr. Tariq regularly attends regional and international congresses to share his vast experience with colleagues and to evaluate and introduce proven and successful new treatment regimes for his patients.

Dr. Tariq's team consists of fully qualified plastic surgeons and dermatologists who have undergone rigorous training in Plastic, Reconstructive Surgery and Dermatology in addition to their continuous training and upgrading of skills under the direction of Dr. Tariq himself.

Dr. Tariq and his team have extensive experience in performing the aesthetic procedures explained in this booklet. If you decide on surgery, you can be assured that our entire staff will provide you with the highest quality medical care.

A WORD ABOUT BREAST REDUCTION IN MEN...

Gynaecomastia is a medical term that comes from the Greek words for “women-like breasts”. Though this oddly named condition is rarely talked about, it’s actually quite common. Gynaecomastia affects an estimated 40 to 60 percent of men. It may affect only one breast or both. Though certain drugs and medical problems have been linked with male breast overdevelopment, there is no known cause in the vast majority of cases.

For men who feel self-conscious about their appearance, breast-reduction surgery can help. The procedure removes fat and or glandular tissue from the breasts, and in extreme cases removes excess skin, resulting in a chest that is flatter, firmer, and better contoured.

If you’re considering surgery to correct gynaecomastia, this brochure will give you a basic understanding of the procedure when it can help, how it’s performed, and what results you can expect. It can’t answer all of your questions, since a lot depends on your individual circumstances. Please be sure to ask your doctor if there is anything about the procedure you don’t understand.

THE BEST CANDIDATES FOR GYNAECOMASTIA CORRECTION

Surgery to correct Gynaecomastia can be performed on healthy, emotionally stable men of any age. The best candidates for surgery have firm, elastic skin that will reshape to the body's new contours.

Surgery may be discouraged for obese men, or for overweight men who have not first attempted to correct the problem with exercise or weight loss. Also, individuals who drink alcoholic beverages in excess or smoke drugs are usually not considered good candidates for surgery. Drugs, along with anabolic steroids, may cause gynaecomastia. Therefore, patients are first directed to stop the use of these drugs to see if the breast fullness will diminish before surgery is considered an option.

ALL SURGERY CARRIES SOME UNCERTAINTY AND RISK


When male breast-reduction surgery is performed by a qualified plastic surgeon, complications are infrequent and usually minor. Nevertheless, as with any surgery, there are risks. These include infection, skin injury, excessive bleeding, adverse reaction to anaesthesia, and excessive fluid loss or accumulation. The procedure may also result in noticeable scars, permanent pigment changes in the breast area, or slightly mismatched breasts or nipples. If asymmetry is significant, a second procedure may be performed to remove additional tissue.

The temporary effects of breast reduction include loss of breast sensation or numbness, which may last up to a year.

PLANNING YOUR SURGERY

The initial consultation with your surgeon is very important. Your surgeon will need a complete medical history, so check your own records ahead of time and be ready to provide this information. First, your surgeon will examine your breasts and check for causes of the gynaecomastia, such as impaired liver function, use of estrogen-containing medications, or anabolic steroids. If a medical problem is the suspected cause, you'll be referred to an appropriate specialist.

Your plastic surgeon may, in extreme cases, also recommend a mammogram, or breast x-ray. This will not only rule out the very small possibility of breast cancer, but will reveal the breast's composition. Once your surgeon knows how much fat and glandular tissue is contained within the breasts, he or she can choose a surgical approach to best suit your needs.



Many men have gynaecomastia enlarged, female-like breasts caused by excess glandular tissue or fat (or both).

Don't hesitate to ask your surgeon any questions you may have during the initial consultation- including your concerns about the recommended treatment or the costs involved. Treatment of gynaecomastia may be covered by medical insurance, but policies vary greatly. Check your policy or call your insurance company to be sure. If you are covered, make certain you get written pre-authorization for the treatment recommended by your surgeon.

PREPARING FOR YOUR SURGERY

Your surgeon will give you specific instructions on how to prepare for surgery, including guidelines on eating, drinking, and taking certain vitamins and medications.

Smokers should plan to stop smoking for a minimum of one to two weeks before surgery and during recovery. Smoking decreases circulation and interferes with proper healing. Therefore, it is essential to follow all your surgeon's instructions.

WHERE YOUR SURGERY WILL BE PERFORMED

Surgery for gynaecomastia may be performed as an outpatient procedure, but in many cases, or those where other medical conditions present cause for concern, an overnight hospital stay may be recommended. The surgery itself usually takes about an hour and a half to complete. However, more extensive procedures may take longer.

TYPE OF ANAESTHESIA

Correction of enlarged male breasts may be performed under general, or in some cases, under local anaesthesia plus sedation. You'll be awake, but very relaxed and insensitive to pain. More extensive correction may be performed under general anaesthesia, which allows the patient to sleep through the entire operation. Your surgeon will discuss which option is recommended for you, and why this is the option of choice.

THE SURGERY

If excess glandular tissue is the primary cause of the breast enlargement, it will be excised, or cut out, with a scalpel. The excision may be performed alone or in conjunction with liposuction. In a typical procedure, an incision is made in an inconspicuous location, either on the edge of the areola or in the



Glandular tissue must be cut out, usually through a small incision near the edge of the areola.

underarm area. Working through the incision, the surgeon cuts away the excess glandular tissue, fat and skin from around the areola and from the sides and bottom of the breast. Major reductions that involve the removal of a significant amount of tissue and skin may require larger incisions that result in more conspicuous scars. If liposuction is used to remove excess fat, the cannula is usually inserted through the separate incisions.

If your gynaecomastia consists primarily of excessive fatty tissue, your surgeon will likely use liposuction to remove the excess fat. A small incision, less than a half-inch in length, is made around the edge of the areola, the dark skin that surrounds the nipple, or the incision may be placed in the underarm area. A slim hollow tube called a cannula which is attached to a vacuum pump, is then inserted into the incision. Using strong, deliberate strokes, the surgeon moves the cannula through the layers beneath the skin, breaking up the fat and suctioning it out. Patients may feel a vibration or some friction during the procedure, but generally no pain.



Fatty tissue can be removed by liposuction. A small, hollow tube is inserted through a tiny incision, leaving a nearly imperceptible scar.

In extreme cases where large amounts of fat or glandular tissue have been removed, skin may not adjust well to the new smaller breast contour. In these cases, excess skin may have to be removed to allow the remaining skin to firmly re-adjust to the new breast contour.

Sometimes, a small drain is inserted through a separate incision to draw off excess fluids. Once closed, the incisions are usually covered with a dressing. The chest may be wrapped to keep the skin firmly in place.

AFTER YOUR SURGERY

Whether you've had excision with a scalpel or liposuction, you will feel some discomfort for a few days after surgery. However, discomfort can be controlled with medications prescribed by your surgeon. In any case, you should arrange to have someone drive you home after surgery and to help you out for a day or two if needed.

You'll be swollen and bruised for a while in fact, you may wonder if there's been any improvement at all. To help reduce swelling, you'll probably be instructed to wear an elastic pressure garment continuously for a week or two, and for a few weeks longer at night. Although the worst of your swelling will dissipate in the first few weeks, it may be three months or more before the final results of your surgery are apparent.

In the meantime, it is important to begin getting back to normal. You'll be encouraged to begin walking around on the day of surgery, and can return to work when you feel well enough which could be as early as a day or two after surgery. Any stitches will generally be removed about 1 to 2 weeks following the procedure.


Your surgeon may advise you to avoid heavy exercise for about six weeks. You'll be told to stay away from any sport or job that risks a blow to the chest area for at least four weeks. In general, it will take about a month before you're back to all of your normal activities.

You should also avoid exposing the resulting scars to the sun for at least six months. Sunlight can permanently affect the skin's pigmentation, causing the scar to turn dark. If sun exposure is unavoidable, use a strong sunblock.


YOUR NEW LOOK

Gynaecomastia surgery can enhance your appearance and self-confidence, but it won't necessarily change your looks to match your ideal. Before you decide to have surgery, think carefully about your expectations and discuss them frankly with your plastic surgeon.

The results of the procedure are significant and permanent. If your expectations are realistic, chances are good that you'll be very satisfied with your new look.



Following surgery for gynaecomastia, the patient has a more masculine chest contour.



Gynaecomastia

Appointments المواعيد

Telephone: +973 17 822822 هاتف:
Laser Centre: +973 17 822820 مركز الليزر:
+973 17 822821
Fax: +973 17 822899 فاكس:
SMS: +973 39108828 SMS:
e-mail: tariplas@batelco.com.bh البريد الإلكتروني:
website: www.dermaplast.com.bh موقع الإنترنت:

To make an appointment للحصول على موعد الرجاء
please contact the above الإتصال بالأرقام الموضحة
numbers. أعلاه.

If you must cancel, we إذا أردت إلغاء الموعد
ask that you please الرجاء إخبارنا وذلك
notify us by telephone, بالإتصال هاتفياً أو إرسال
sms or e-mail as early as رسالة قصيرة أو عبر
possible. البريد الإلكتروني.



درما بلاست
dermaplast

مستشفى الدكتور طارق
DR. TARIQ HOSPITAL

ص.ب: ٣٢٢٢٤، مبنى: ٢٨٤، طريق: ٣٣٠٦، مجمع: ٣٣٢، الماحوز، مملكة البحرين
P.O.Box: 32234, Building: 284, Road: 3306, Block: 333
Mahooz, Kingdom of Bahrain